

(2)

AD-A211 289

REPORT DOCUMENTATION PAGE

CTE
4 1989

D

1b. RESTRICTIVE MARKINGS

THE FILE COPY

2b. DECLASSIFICATION/DOWNGRADING SCHEDULE

3. DISTRIBUTION/AVAILABILITY OF REPORT

Approved for public release;
distribution unlimited.

4. PERFORMING ORGANIZATION REPORT NUMBER

5. MONITORING ORGANIZATION REPORT NUMBER(S)

ARO 23767.7-LS-F

6a. NAME OF PERFORMING ORGANIZATION

North Carolina State University

6b. OFFICE SYMBOL
(If applicable)

7a. NAME OF MONITORING ORGANIZATION

U. S. Army Research Office

6c. ADDRESS (City, State, and ZIP Code)

Raleigh, N.C. 27695-7612

7b. ADDRESS (City, State, and ZIP Code)

P. O. Box 12211
Research Triangle Park, NC 27709-22118a. NAME OF FUNDING/SPONSORING
ORGANIZATION

U. S. Army Research Office

8b. OFFICE SYMBOL
(If applicable)

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

DAA403-86-G-0035

8c. ADDRESS (City, State, and ZIP Code)

P. O. Box 12211
Research Triangle Park, NC 27709-2211

10. SOURCE OF FUNDING NUMBERS

PROGRAM
ELEMENT NOPROJECT
NOTASK
NOWORK UNIT
ACCESSION NO

11. TITLE (Include Security Classification)

Membrane Fusion: The Role of Polyphosphatidylinositol

12. PERSONAL AUTHOR(S)
Wendy F. Boss

Jeffery J. Wheeler

13a. TYPE OF REPORT
Final13b. TIME COVERED
FROM 3-86 TO 3-8914. DATE OF REPORT (Year, Month, Day)
89-7-2415. PAGE COUNT
3

16. SUPPLEMENTARY NOTATION

The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

17. COSATI CODES

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

FIELD GROUP SUB-GROUP

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

We have found that the presence of the phospholipid, lysophosphatidylinositol monophosphate (LPIP) correlates positively with the fusion potential of fusogenic carrot cells. There was no correlation with the presence of phosphatidylinositol monophosphate (PIP) and phosphatidylinositol biphosphate (PIP₂) and fusion. Nor was there evidence for the need for PIP or PIP₂ turnover in order for the cells to be fusion permissive. LPIP was synthesized primarily from the phosphorylation of lysophosphatidylinositol. Lysolipids were found to decrease the phosphorylation of PI and PIP suggesting a mechanism for regulating the biosynthesis of the polyphosphoinositides which are key components of the signal transduction pathway in many animal cells.

89

3

11

005

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT

☐ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION

Unclassified

22a. NAME OF RESPONSIBLE INDIVIDUAL

22b. TELEPHONE (Include Area Code)

22c. OFFICE SYMBOL

MEMBRANE FUSION: THE ROLE OF POLYPHOSPHATIDYLINOSITOL

FINAL REPORT

Wendy F. Boss and Jeffery J. Wheeler

U. S. Army Research Office

Grant Number: DAAL03-86-G-0035

North Carolina State University

Approved for Public Release: Distribution Unlimited

Brief Outline of Research Findings

The focus of this research was to investigate the role of the polyphosphoinositides in cell fusion. As a result of our research we have found that the presence of the phospholipid, lysophosphatidylinositol monophosphate (LPIP) correlates positively with the fusion potential of fusogenic carrot cells. There was no correlation with the presence of phosphatidylinositol monophosphate (PIP) and phosphatidylinositol bisphosphate (PIP₂) and fusion. Nor was there evidence for the need for PIP or PIP₂ turnover in order for the cells to be fusion permissive.

Although LPIP was a known lysophospholipid, it had not been reported in a living system prior to our work. For this reason we were careful to identify the lipid and have used fast atom bombardment mass spectrometry to identify the fatty acids associated with the lipid.

We also found that LPIP was synthesized by ATP-dependent phosphorylation of LPI. While LPI was found primarily in the intracellular membranes LPIP was found predominantly in the plasma membrane. Kinase activity, however, was found in both membrane fractions in the in vitro assay system. Thus the question remained as to what regulated the biogenesis of the plasma membrane LPIP in vivo. This has been addressed to some extent in a paper recently submitted for publication. Conditions which favor formation of LPI from phospholipase A₂ hydrolysis of PI (acidic pH, pH 5.5) do not favor kinase activity. Thus our working hypothesis is that the LPI is synthesized on the intracellular membranes in an acidic secretory vesicle and transported to the plasma membrane where it is phosphorylated.

In addition, we found that the lysolipids affect the biosynthesis of PIP and PIP₂ which are involved in signal transduction in many animal cells. Thus the lysolipid pathway may be important for regulating cellular signalling as well as membrane fusion.

We were unable to do biophysical studies using spin labeled polyphosphoinositides due to time constraints and the difficulty in synthesizing derivatives of these lipids. To my knowledge there are no published reports of spin label phosphoinositides, although Dr. S. McLaughlin says his lab has recently been successful in synthesizing a fluorescent analogue.

During the three years of the fellowship, Dr. Wheeler has had two manuscripts published in journals, co-authored two book chapters (first author of one) and has two manuscripts submitted for publication. These are listed below and a copy of the latest manuscripts is attached.

We feel very fortunate to have had support from the Army Research

Office to pursue this work. While we have gained new information about the structure of a naturally fusing cell membrane and the biosynthesis of lysophosphoinositides, we have produced a fine scientist who has had offers for postdoc positions from excellent laboratories. Dr. Wheeler will be joining either the laboratory of Dr. Pieter Cullis or Dr. R. M. Epand to pursue biophysical studies of fusion. Thank you for your support.

Publications:

Wheeler, JJ and WF Boss. 1987. Polyphosphoinositides are present in the plasma membranes isolated from fusogenic carrot cells. Plant Physiol. 85: 389-392.

Boss, WF and JJ Wheeler. 1989. Strategies for optimizing cell fusion. In: Biotechnology Vol 7b. Verlagsgesellschaft. West Germany.

Wheeler JJ and WF Boss. 1989. The presence of sn-1-palmitoyl phosphatidylinositol monophosphate correlates positively with the fusogenic state of the plasma membrane of fusogenic carrot cells grown in suspension culture. Biochim Biophys Acta. (in press).

Wheeler, JJ and Wf Boss. 1989. Inositol Lysolipids. In: Inositol Metabolism in Plants. DJ Morre, WF Boss, FA Loewus (eds) Alan R. Liss, New York. (in press).

Van Breemen, RB, JJ Wheeler, WF Boss. Identification of Carrot Inositol Phospholipids by Fast Atom Bombardment Mass Spectrometry. Lipids (submitted).

Wheeler, JJ and WF Boss. Phosphorylation of LPI by plasma membranes isolated from carrot cells. Acta Biochem Biophys (submitted).

Degrees awarded: Ph. D. Jeffery J. Wheeler. May, 1989

Submitted By: Wendy F. Boss
North Carolina State University
Box 7612
Raleigh, NC 27695
919-737-3496



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	